

# DECCO



## DECCO PENBOTEC® 400 SC

For postharvest use on pome fruits

*Active Ingredient:*

Pyrimthanil: 4,6-dimethyl-N-phenyl-2-pyrimidinamine\* 37.14%

*Other Ingredients:* 62.86%

*Total:* 100.00%

Contains 3.27 lbs. active ingredient per gallon (400 g/L)

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

FIRST AID	
<b>If on skin</b>	-Take off contaminated clothing. -Rinse skin immediately with plenty of water for 15-20 minutes. -Call a poison control center or doctor for treatment advice.
<b>If inhaled</b>	-Move person to fresh air. -If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. -Call a poison control center or doctor for further treatment advice
<b>If swallowed</b>	-Call a poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to do so by a poison control center or doctor. -Do not give anything by mouth to an unconscious person.
<b>HOT LINE NUMBER:</b> For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. For chemical emergency assistance (spill, leak, fire, or accident), call Chem Trec at 1-800 424-9300.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

NET CONTENTS: 33.8 fl. oz. (1 liter)

EPA REG. NO. 43813-32-2792

EPA EST. No.: 074263-DEU-001; 43813-001; 43813-002

**Decco Cerexagri Inc.**  
**1713 S. California Ave.**  
**Monrovia, CA 91016-0120**

## **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS**

### **CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A). Remove and wash contaminated clothing before reuse. Avoid breathing spray mist.

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **STORAGE AND DISPOSAL**

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### **GENERAL INFORMATION:**

PENBOTEC 400 SC fungicide controls blue and gray mold on pome fruit. Use of PENBOTEC 400 SC should be integrated into an overall disease management program within each packinghouse.

## FUNGICIDE RESISTANCE MANAGEMENT

### GROUP 9 FUNGICIDE

PENBOTEC 400 SC is an anilinopyrimidine fungicide that does not exhibit cross-resistance to sterol-inhibiting or benzimidazole fungicides. PENBOTEC 400 SC interferes with the secretion of enzymes necessary for infection by several postharvest pathogens, therefore, preventative applications soon after harvest or injury is necessary for optimal product performance. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of postharvest resistance management strategies should be practiced. These may include rotating and/or tank mixing with products having different modes of action. Responsible resistance management practices are necessary to ensure the long-term effectiveness for decay control.

#### MIXING INSTRUCTIONS

Fill tank with half of the required amount of water or water wax and start mechanical agitation. Add the required amount of PENBOTEC 400 SC and then add the remaining volume of water or water wax. Maintain agitation after mixing and do not allow treating solution to stand overnight or for prolonged periods. For in-line injection systems better distribution will be achieved by injecting a larger volume of a more dilute solution per unit of time.

The physical compatibility of PENBOTEC 400 SC with all tank mix partners and wax types has not been fully investigated. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rates for the individual products may be exceeded. Prior to tank mixing, conduct a jar test with the volumes and rates typically used for postharvest disease control. Add proportionate amounts of the products to a small container of water or water wax in the following order: Wettable powders and water-dispersible granules first, followed by liquid flowables, and emulsifiable concentrates last. After thorough mixing, let stand for at least 15 minutes. Separation, globules, sludge, flakes or other precipitates are indicative of physical incompatibility. Physical compatibility is indicated if the combination remains mixed or can be remixed readily. THE CROP SAFETY OF ALL POTENTIAL PESTICIDE TANK MIXES WITH PENBOTEC 400 SC INCLUDING ANTI-SCALD AGENTS AND WAXES, HAS NOT BEEN TESTED ON ALL CROPS AND VARIETIES. BEFORE APPLYING ANY TANK MIXTURE, SAFETY TO THE TARGET CROP SHOULD BE CONFIRMED.

#### USE RATES:

<i>POME FRUITS</i>			
Treatment Method	Rate Fl.oz/100 gal	Treatment Time (min)	Concentration A.I. (ppm)
Dip, wash tanks	16 – 32	1	500-1000
Drenchers	16 - 32	1	500-1000
Aqueous line spray	32	1	1000
Wax line spray	64	1	2000

Periodically measure the PENBOTEC 400 SC concentration of the treating solutions from drenchers, dips and wash tanks to maintain the proper level of active ingredient. Fruits treated by aqueous or wax line spray applications should be cleaned prior to application. Aqueous applications must be applied prior to fruit prior to receiving any wax applications.

### Treatment Schemes

Double treatment with PENBOTEC 400 SC is the maximum allowed on pome fruits. Acceptable multiple treatment scenarios are presented below:

Treatment Scenario	PENBOTEC 400 SC Application Methods			
	Drench	Dip & Wash tanks	Aqueous spray	Shipping Wax
<b>POME FRUIT</b>				
1	√	√		
2	√		√	
3	√			√
4		√		√
5			√	√
6		√	√	

### WARRANTY AND DISCLAIMER

CEREXAGRI INC. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. CEREXAGRI INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT ALLOWED BY APPLICABLE LAW, IN NO CASE SHALL CEREXAGRI INC OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST, OR OTHER EXPENSES INCURRED IN REPACKAGING, SORTING OR REPROCESSING.

CEREXAGRI INC. and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of CEREXAGRI INC

**DECCO**  
**Cerexagri Inc.**  
**1713 S. California Ave.**  
**Monrovia, CA 91017-0120**

® Registered trademark of Janssen Pharmaceutica N.V.

\* Pyrimethanil manufactured by Bayer CropScience AG

Rev. 04/05